

COMPUTEPREP

TIMED CALCULATION DRILLS · BANK & SSC EDITION

Free Sample: 23 Timed Speed Drills (One of Every Type)

Free sample — 23 engine-minted drills, one of every drill type ComputePrep generates, across Easy, Medium & Hard. Same chapters, same orientation cards, same exam-pace target times and same worked answer key as the full 150-drill Speed Book — just one drill per type instead of a full practice set. Free to keep and free to forward.

Sectioned by Family

All 23 Drill Types

Exam-Pace Target Times

Free Sample · 2026 · Share It Freely

How to Use This Book

Every drill is a multiple-choice question with one defensible answer and a **target time** printed under it. That target is not editorial — it is the exact per-question clock the ComputePrep daily enforces for that drill type at that difficulty.

Work against the clock, not just for the answer. A quant paper is not decided by whether you *can* do a sum; it is decided by whether you can do it inside the time the paper allows. Getting a drill right in triple the target time is a miss, and this book is written so you can see that happening.

Method: start a timer, work it mentally wherever the technique allows, commit to an option before you check, then read the worked solution even when you were right — the method in the answer key is often shorter than the one you used, and the shorter method is the one that survives exam pressure. Re-drill any type you missed on *time*, not just the ones you got wrong.

Start with Chapter 1. Speed & Vedic Shortcuts is deliberately the largest chapter: percentages, profit & loss, interest, averages, DI and mensuration all bottom out in a multiplication or a square, so every second you take off those makes the rest of the book — and the paper — faster.

Every answer here is **engine-computed, never hand-written** — each drill is minted by the same pure engine that mints the ComputePrep daily, and each worked solution is that engine's own step-by-step working. The distractors are built from the specific arithmetic slips the technique invites (a dropped carry, a base confusion, a percentage applied to the wrong number), so picking one tells you exactly which step went wrong.

The First 10 Seconds of Any Calculation

Before you compute anything, run this checklist — the ten seconds a fast solver spends on every question, whether it is a multiplication, a percentage, a DI cell or a mensuration formula. Speed in a quant paper is not faster arithmetic; it is less arithmetic.

1. **Look at the numbers before the operation.** Ending in 5? Both near 100? Same leading digits with units adding to 10? Close together, so their midpoint is easy to square? Each of those is a named technique in Chapter 1 and turns the whole multiplication into one step.
2. **Round, then correct.** 98×47 is 100×47 minus 2×47 . Most "hard" multiplications are an easy one plus a small correction — find the easy one first.
3. **Check what the question actually asks.** Approximate value? Percentage *change* or the new value? The profit or the selling price? More marks are lost here than to arithmetic.
4. **Scan the options before you compute.** They are often decades apart, or only one is even/odd, or only one is in range — in which case an estimate finishes the question and the exact working is wasted time.
5. **Estimate the magnitude first.** Know roughly where the answer must land, then compute. It costs a second and catches the misplaced decimal or the dropped zero before you commit.
6. **Decide fast whether to skip.** If nothing above fires within ten seconds, this is a long-working question. Mark it, move on, come back. Protecting the clock is a skill you drill here too.
7. **Never write down what you can carry.** Each intermediate you write is time and a copying error. The techniques in Chapter 1 are built so the intermediate fits in your head.

Contents

23 drills in 5 sections, plus a worked solution for every one of them. Drills are numbered **1–23 straight through the book** — the section lists that follow are slices of that one run, not separate books, and the answer key at the back uses the same numbers.

1 Speed & Vedic Shortcuts

The calculations that decide a quant section — squares, products, and long BODMAS chains — done by pattern instead of by long multiplication.

Drills 1–10

10 drills · 4E / 3M / 3H

2 Percentages & Commerce

Percentage, profit, interest, and averages — one family of problems, and every one of them turns on knowing which number the percentage is charged against.

Drills 11–14

4 drills · 1E / 2M / 1H

3 Time, Work & Motion

Work done, distance covered, and a stream helping or fighting you — all three are rate problems, and rates add where times never do.

Drills 15–17

3 drills · 1E / 1M / 1H

4 Data Interpretation & Equations

Tables, caselets, and paired quadratics — the marks are in reading only what the question asks for, then computing once.

Drills 18–20

3 drills · 1E / 1M / 1H

5 Counting & Measurement

Arrangements, chances, and shapes — the topics where the setup is the whole battle and the arithmetic afterwards is trivial.

Drills 21–23

3 drills · 1E / 1M / 1H



Solutions

The answer plus the full worked method for every drill, in the same numbering.

All 23

answer key

How to move around: every section opens with a one-page guide to its drill types, then its own contents list; each drill header carries a link back to that list. Clues are numbered — if something looks wrong, write to hello@computeprep.com quoting the drill number and the clue number (e.g. "Drill 42, clue 3"). More at computeprep.com.

Speed & Vedic Shortcuts — How to Read This Section

The calculations that decide a quant section — squares, products, and long BODMAS chains — done by pattern instead of by long multiplication.

Speed Arithmetic. ONE operation on exam-sized numbers, against a countdown — an addition, a subtraction, a multiplication, or (from medium up) a division. There is no method to choose here; the whole skill is doing a single step accurately without reaching for paper, which is what the rest of this chapter then builds on.

WORKED EXAMPLE $34 \times 26 = ?$ Split the second factor instead of stacking it: $34 \times 20 = 680$, and $34 \times 6 = 204$. $680 + 204 = 884$.

COMMON MISTAKE checking your digits but not your MAGNITUDE. The wrong options are planted close on purpose — one out at the units, one out by ten, one out by a hundred — so an answer that merely looks about right is exactly the one that was left there for you.

Squares, Cubes & Roots. Squares, cubes and their roots for the numbers the exam actually uses. Two facts do most of the work: a square's LAST DIGIT is fixed by the root's last digit, and the nearest round square brackets the answer.

WORKED EXAMPLE $\sqrt{5184} = ?$ It ends in 4, so the root ends in 2 or 8; and $70^2 = 4,900$ while $80^2 = 6,400$, so the root is in the 70s. $72^2 = 5,184$, so the root is 72.

COMMON MISTAKE guessing from the options without the last-digit test — that one check alone kills most distractors in about a second.

Complementary Products. When two numbers share the same LEADING part AND their units digits add to 10, the product assembles itself (Antyayordashake): the lead $\times$ its successor gives the front, the two units multiplied give the last two digits. It works at any width — 245×245 is the same trick on the lead 24.

WORKED EXAMPLE $43 \times 47 = ?$ Same lead 4, and $3 + 7 = 10$. Front: $4 \times 5 = 20$. Tail: $3 \times 7 = 21$. $= 2,021$.

COMMON MISTAKE forgetting the tail is always TWO digits — 91×99 gives front $9 \times 10 = 90$ and tail $1 \times 9 = 9$, which must be written 09, so 9,009 and not 909.

Simplification (BODMAS). One expression, one answer, and everything turns on ORDER: brackets first, then division and multiplication together left to right, then addition and subtraction together left to right. Two operations at easy; four, with a bracket in front, by the time you reach hard.

WORKED EXAMPLE $83 + 9 \times 9 - 70 \div 7 = ?$ $\times$ and $\div$ outrank $+$ and $-$, so clear those first: $9 \times 9 = 81$ and $70 \div 7 = 10$. Now $83 + 81 - 10 = 154$.

COMMON MISTAKE working strictly left to right — doing the $83 + 9$ before the multiplication that outranks it. It is the one slip this kind exists to catch, and it always lands you on a number that looks perfectly reasonable.

Vedic Squaring. Two sutras for squares the exam loves. Numbers ending in 5 (Ekadhikena): multiply the head by its successor and append 25. Numbers near a base like 100 (Yavadunam): slide by the deficit and append the deficit's square.

WORKED EXAMPLE $85^2 = ?$ It ends in 5, so take the head 8 and its successor 9: $8 \times 9 = 72$, then append 25. $= 7,225$.

COMMON MISTAKE using the head $\times$ successor trick on a number that does NOT end in 5 — it is a rule about the units digit, not a general squaring shortcut.

Difference of Squares. Any two numbers multiply to the square of their MIDPOINT minus the square of the distance from it (Sankalana-vyavakalanabhyam). The midpoint is just their average, and when the two sit close together that distance is 1, 2 or 3 — so one square replaces a whole multiplication.

WORKED EXAMPLE $67 \times 73 = ?$ Their midpoint is 70 and both sit 3 away, so the product is $70^2 - 3^2$. $4,900 - 9 = 4,891$.

COMMON MISTAKE subtracting the distance instead of its SQUARE — writing $4,900 - 3$. The identity is $(m - d)(m + d) = m^2 - d^2$, and the d is squared.

Nikhilam. Multiplying two numbers near a base (98×93 , 104×107): work with their small deficits or surpluses instead of the numbers themselves. Cross-subtract (or cross-add) for the head; multiply the two deviations for the tail.

WORKED EXAMPLE $98 \times 93 = ?$ Deficits from 100 are 2 and 7. Head: $98 - 7 = 91$ ($93 - 2$ gives the same). Tail: $2 \times 7 = 14$. = 9,114.

COMMON MISTAKE letting the tail overflow its two digits without carrying — 87×89 has deficits 13 and 11, so the tail is 143: carry the 1 into the head ($76 + 1 = 77$) and keep 43, giving 7,743.

Paravartya Division. Paravartya Yojayet, for dividing by 11: the flag digit ($11 = 10 + 1$, so the flag is 1) gets NEGATED, then chains leftward — bring down the first digit as-is, and each next quotient digit is the dividend digit plus ($-1 \times$ the previous quotient digit). The last digit falls out the same way and gives the remainder for free.

WORKED EXAMPLE $475 \div 11 = ?$ Flag digit is 1 ($11 = 10 + 1$); negate it to -1 . Bring down 4, then $7 + (-1 \times 4) = 3$. Quotient 43, remainder 2 — check: $43 \times 11 + 2 = 475$.

COMMON MISTAKE ADDING the previous quotient digit instead of subtracting it — forgetting to negate the flag digit. The slip doesn't stay local: it cascades, so every digit after the first comes out wrong, not just one.

Crosswise Multiplication. Urdhva-Tiryagbhyam, the general-purpose one: a multiplication in a single sweep — vertical on the left, crosswise in the middle, vertical on the right — with nothing written down but the answer. The easy drills are its $\times 11$ special case (add each pair of neighbours); the hard ones run three digits by two.

WORKED EXAMPLE $23 \times 41 = ?$ Vertical left: $2 \times 4 = 8$. Crosswise: $2 \times 1 + 3 \times 4 = 14$. Vertical right: $3 \times 1 = 3$. Lay them out as $8 \mid 14 \mid 3$. The middle column carries its ten into the front: 943.

COMMON MISTAKE writing the three column results side by side without carrying — $8 \mid 14 \mid 3$ read straight off gives 8,143 instead of 943.

Anurupyena Multiplication. Multiplying two numbers close to a working base that ISN'T a power of ten — 50, since $50 = 100 \div 2$. Cross-add one number with the other's deviation for a raw left part, then RESCALE it by the base's ratio to 100 (halve it, for base 50). The deviations' own product is the right part, padded to two digits.

WORKED EXAMPLE $54 \times 52 = ?$ Deviations from 50: +4 and +2. Cross-add: $54 + 2 = 56$, then rescale by the base's ratio to 100 ($\div 2$): $56 \div 2 = 28$. Right part: $4 \times 2 = 08$. Join them: $28 \mid 08 \rightarrow 2,808$.

COMMON MISTAKE joining the raw cross-add without rescaling it — writing $56 \mid 08$ straight as 5,608. The working base is 50, not 100, so that left part is always twice too big until you divide it by the base's ratio.

Section 1 Contents — Speed & Vedic Shortcuts

Drills 1–10 of this section only. The full book index is at the front.

Drill 1 · Easy

Drill 2 · Medium

Drill 3 · Hard

Drill 4 · Easy

Drill 5 · Medium

Drill 6 · Hard

Drill 7 · Easy

Drill 8 · Medium

Drill 9 · Hard

Drill 10 · Easy

Drill 1 EASY *Speed Arithmetic*[↑ Section index](#)

Solve in your head — beat the clock.

Given:

$$8 \times 9 = ?$$

Options:

- A** 72
- B** 81
- C** 84
- D** 80
- E** 82

Target time: ~15s

⚠ **Common mistake:** 8×9 is one step, so the slip is a table error, not a method error — $8 \times 10 = 80$ and $8 \times 8 = 64$ are both on the board.

Drill 2 MEDIUM *Simplification (BODMAS)*[↑ Section index](#)

Simplify — order of operations wins.

Given:

$$71 + 4 \times 4 - 45 \div 9 = ?$$

Options:

- A** 92
- B** 82
- C** 81
- D** 72
- E** 28

Target time: ~30s

⚠ **Common mistake:** BOTH the $\times$ and the $\div$ resolve before any $+$ or $-$: settle $4 \times 4 = 16$ and $45 \div 9 = 5$ first, then walk the additions. Letting the $\div 9$ swallow the whole expression gives 28.

Drill 3 HARD *Squares, Cubes & Roots*[↑ Section index](#)

Powers and roots — instant recall.

Given:

$$\sqrt[3]{343} = ?$$

Options:

- A** 6
- B** 5
- C** 7
- D** 0
- E** 8

Target time: ~30s

⚠ **Common mistake:** Cubes make the last digit unique: 343 ends in 3, so the root ends in 7 — no other digit does that. The magnitude pins the rest, so there is nothing to estimate.

Drill 4 EASY *Vedic Squaring*[↑ Section index](#)

Squaring a number ending in 5 — beat the clock.

Given:

$$55^2 = ?$$

Options:

- A** 3014
- B** 3016
- C** 3625
- D** 3025
- E** 2525

Target time: ~15s

⚠ **Common mistake:** Ekadhikena multiplies the head by ONE MORE than itself — 5×6 , not 5×5 . The tail is always exactly 25.

Drill 5 MEDIUM *Crosswise Multiplication*[↑ Section index](#)

Crosswise multiplication — beat the clock.

Given:

$$63 \times 84 = ?$$

Options:

- A 5282
- B 5052
- C 5292
- D 5287
- E 4882

Target time: ~30s

⚠ **Common mistake:** The middle band takes BOTH cross-products — $6 \times 4 + 3 \times 8 = 48$, not just one of them, and 48 overflows one digit, so it must carry into the tens.

Drill 6 HARD *Nikhilam*[↑ Section index](#)

Nikhilam multiplication — beat the clock.

Given:

$$106 \times 112 = ?$$

Options:

- A 11800
- B 11306
- C 11872
- D 72
- E 10712

Target time: ~35s

⚠ **Common mistake:** Cross-add ONCE ($106 + 12$) — the two cross-steps give the same left part, so doing both is a wasted step — then pad the tail to exactly 2 digits.

Drill 7 EASY *Complementary Products*[↑ Section index](#)

Units sum to 10, same lead — beat the clock.

Given:

$$88 \times 82 = ?$$

Options:

- A** 7221
- B** 7226
- C** 6416
- D** 7210
- E** 7216

Target time: ~15s

⚠ **Common mistake:** The left part is the head times ONE MORE than itself (8×9), and the right part is a PRODUCT of the units (8×2), always written in two digits.

Drill 8 MEDIUM *Difference of Squares*[↑ Section index](#)

Multiply via the midpoint — beat the clock.

Given:

$$22 \times 38 = ?$$

Options:

- A** 484
- B** 866
- C** 964
- D** 821
- E** 836

Target time: ~20s

⚠ **Common mistake:** Both operands sit 8 either side of 30, so the answer is 30^2 MINUS 8^2 — subtract the small square, never add it, and never stop at 30^2 alone.

Drill 9 **HARD** *Paravartya Division*[↑ Section index](#)

Dividing by 11 — beat the clock.

Given:

$$99055 \div 11 = ?$$

Options:

- A** 9905
- B** 9005
- C** 8992
- D** 9883
- E** 90050

Target time: ~45s

⚠ **Common mistake:** The flag digit is NEGATED — each step subtracts the previous quotient digit, it does not add it. Adding instead cascades the error through every digit that follows.

Drill 10 **EASY** *Anurupyena Multiplication*[↑ Section index](#)

Multiplying near a working base — beat the clock.

Given:

$$48 \times 46 = ?$$

Options:

- A** 4408
- B** 8808
- C** 2219
- D** 2200
- E** 2208

Target time: ~15s

⚠ **Common mistake:** The working base is $50 = 100 \div 2$, so the cross-added left part must be HALVED ($44 \div 2 = 22$) before the deviation product is appended. Skipping the rescale, or doubling instead, is the only mistake this method has.

End of Section 1 — Speed & Vedic Shortcuts

You have worked drills 1–10 (10 of drills in this book). Any blank space below this line is deliberate — each section begins on a fresh page. Next: Section 2 — Percentages & Commerce, starting overleaf.

Percentages & Commerce — How to Read This Section

Percentage, profit, interest, and averages — one family of problems, and every one of them turns on knowing which number the percentage is charged against.

Percentages. Three shapes, and this book drills all of them. Most are a straight 'X% of Y' — convert the percentage to the fraction you already know ($10\% = 1/10$, $20\% = 1/5$, $25\% = 1/4$, $50\% = 1/2$, $75\% = 3/4$) and one division finishes it. Some ask the reverse, 'A is what percent of B'. The hardest chain two changes in a row, and there the second is charged on the number the first one left behind.

WORKED EXAMPLE 800 increased by 50%, then decreased by 10%. Final value? $800 \rightarrow 1,200$ after the rise. The 10% is now charged on 1,200, not on 800 — that is 120. $1,200 - 120 = 1,080$.

COMMON MISTAKE charging the second change on the ORIGINAL number, or just netting the two percentages ($+50 - 10 = +40\%$) and landing on 1,120. A percentage is meaningless until you name what it is a percentage of, and after the first change that base has moved.

Simple & Compound Interest. Simple interest is charged on the original principal every year; compound interest is charged on what the principal has grown into. Over 2 years the gap between them is exactly the interest earned on the first year's interest.

WORKED EXAMPLE ₹8,000 for 2 years at 10% p.a. — how much more is CI than SI? $SI = 8,000 \times 10\% \times 2 = 1,600$. $CI = 8,000 \times (1.1^2 - 1) = 8,000 \times 0.21 = 1,680$. CI exceeds SI by ₹80.

COMMON MISTAKE not using the 2-year shortcut as a check: the difference is always $P \times (r/100)^2$, which is ₹80 here. If your CI – SI doesn't match it, the arithmetic slipped.

Profit & Loss. Cost price, marked price, discount, selling price — four numbers and three percentages between them. Profit % is always on COST; discount is always on MARKED PRICE. Getting those two bases right is most of the topic.

WORKED EXAMPLE An article costing ₹250 is marked 40% above cost and sold at a 10% discount. Profit %? Marked price = 350. Selling price = $350 \times 0.9 = 315$. Profit = $315 - 250 = 65$. $65 \div 250 = 26\%$.

COMMON MISTAKE taking the discount off the COST instead of the marked price — every percentage in this topic names its own base, and swapping two of them is the standard trap.

Averages. One relation, $\text{total} = \text{average} \times \text{count}$, asked three ways. Read the total straight off a given mean; recover a MISSING number by subtracting the rest from that total; or handle a REPLACEMENT, where you never rebuild the total at all — the average simply moves by $(\text{change in total}) \div (\text{count})$.

WORKED EXAMPLE The average of 8 numbers is 43. One of them, 54, is replaced by 14. New average? The total falls by $54 - 14 = 40$, spread over the same 8 numbers — that is 5 off the mean. $43 - 5 = 38$.

COMMON MISTAKE rebuilding the whole sum for a replacement question. Only the total's CHANGE matters, so it stays one subtraction and one division however large the numbers get.

Section 2 Contents — Percentages & Commerce

Drills 11–14 of this section only. The full book index is at the front.

Drill 11 · Medium

Drill 12 · Hard

Drill 13 · Easy

Drill 14 · Medium

Drill 11 MEDIUM Percentages[↑ Section index](#)

Percentage drill — no paper.

Given:

A value fell from 44 to 11. By what percent did it change?

Options:

- A 65
- B 85
- C 25
- D 300
- E 75

Target time: ~30s

⚠ **Common mistake:** The base of a percentage change is where it STARTED (44), not where it ended (11). Dividing by 11 gives 300%, and that option is on the board.

Drill 12 HARD Profit & Loss[↑ Section index](#)

QUESTION

What was the cost price?

Given:

An article is sold for ₹8400, at a profit of 5%.

Options:

- A 400
- B 7998
- C 7980
- D 7992
- E 8000

Target time: ~60s

⚠ **Common mistake:** 5% of the SELLING price is ₹420, and that route gives ₹7980 — which is on the option list. Margins are quoted on COST, so treat ₹8400 as 105% and divide.

Drill 13 **EASY** Simple & Compound Interest[↑ Section index](#)**QUESTION****What is the simple interest earned?****Given:****₹10000 is deposited for 2 years at 20% per annum, simple interest.****Options:**

- A** 4000
- B** 4400
- C** 4009
- D** 14000
- E** 4002

Target time: ~40s

⚠ **Common mistake:** Simple interest never earns on itself — ₹2000 in each of the two years, so ₹4000 in total. Doubling nothing and compounding it anyway gives ₹4400, which is offered; ₹2000 alone is one year and is offered too.

Drill 14 **MEDIUM** Averages[↑ Section index](#)

Averages — total and mean, no paper.

Given:**The average of 6 numbers is 50. The other 5 of them add up to 261. What is the remaining number?****Options:**

- A** 39
- B** 50
- C** 250
- D** 261
- E** 300

Target time: ~30s

⚠ **Common mistake:** Rebuild the whole total first ($6 \times 50 = 300$) — the missing number is the GAP between it and 261, not the average 50 and not the total itself. Both of those are on the board.

End of Section 2 — Percentages & Commerce

You have worked drills 11–14 (4 of drills in this book). Any blank space below this line is deliberate — each section begins on a fresh page. Next: Section 3 — Time, Work & Motion, starting overleaf.

Time, Work & Motion — How to Read This Section

Work done, distance covered, and a stream helping or fighting you — all three are rate problems, and rates add where times never do.

Time & Work. Take the whole job as a convenient number of units — the LCM of the given times — so each worker's rate comes out whole. Rates ADD when people work together, and the job divided by the combined rate is the answer.

WORKED EXAMPLE A alone finishes in 12 days, B alone in 18. Together? Call the work 36 units (LCM): A does 3 a day, B does 2 a day, so together 5 a day. $36 \div 5 = 7.2$ days.

COMMON MISTAKE averaging the two times to get 15 days. Rates add; times do not — two workers are always faster than either one alone, so the answer must be below the smaller time.

Boats & Streams. The stream adds to the boat going downstream and subtracts going up. So downstream speed = boat + stream and upstream = boat – stream, which means the boat is the two effective speeds averaged and the stream is half their difference.

WORKED EXAMPLE 30 km downstream in 2 hours and 30 km upstream in 3 hours. Boat and stream speeds?
Downstream = 15 km/h, upstream = 10 km/h. Boat = $(15 + 10) \div 2$, stream = $(15 - 10) \div 2$. Boat 12.5 km/h, stream 2.5 km/h.

COMMON MISTAKE using the still-water speed where the question wants the effective one (or the reverse) — read carefully which of the three speeds is being asked for.

Time, Speed & Distance. One relation, distance = speed $\times$ time, applied to each leg separately. The one thing to check every time is what is held equal between the legs — equal distances and equal times give different answers.

WORKED EXAMPLE Half a journey at 40 km/h and half at 60 km/h. Average speed? The DISTANCES are equal, so the average is the harmonic mean: $2 \times 40 \times 60 \div (40 + 60) = 48$ km/h, not 50.

COMMON MISTAKE averaging the two speeds. That is only right when equal TIMES are spent at each — with equal distances more time is spent at the slower speed, which drags the average down.

Section 3 Contents — Time, Work & Motion

Drills 15–17 of this section only. The full book index is at the front.

Drill 15 · Hard

Drill 17 · Medium

Drill 16 · Easy

Drill 15 **HARD** *Time & Work*[↑ Section index](#)**QUESTION****Working together, in how many days will they finish it?****Given:**

1. Anil can complete a piece of work in 6 days.
2. Bhavna can complete the same work in 12 days.

Options:

- A** 3
- B** 4
- C** 1
- D** 18
- E** 0

Target time: ~70s

⚠ **Common mistake:** Days do not add. $6 + 12 = 18$ days is on the option list and is slower than either of them working alone, which cannot be right — add the RATES ($2 + 1$ units/day) and divide the work by the sum.

Drill 16 **EASY** *Time, Speed & Distance*[↑ Section index](#)**QUESTION****How far does it travel?****Given:****A train runs at a steady 30 km/h for 3 hours.****Options:**

- A** 90
- B** 99
- C** 120
- D** 33
- E** 98

Target time: ~40s

⚠ **Common mistake:** Speed and time MULTIPLY here — 30×3 . The units tell you which way round: $\text{km/h} \times \text{h}$ leaves km, while $30 \div 3 = 10$ (offered) would leave km/h^2 .

Drill 17 MEDIUM Boats & Streams[↑ Section index](#)**QUESTION****How far does it go?****Given:**

1. A boat's speed in still water is 16 km/h and the stream flows at 5 km/h.
2. It travels upstream for 5 hours.

Options:

- A** 55
- B** 46
- C** 105
- D** 11
- E** 41

Target time: ~50s

⚠ **Common mistake:** Going upstream the stream FIGHTS you, so subtract it: $16 - 5 = 11$ km/h. Using 21 km/h (the other direction) or ignoring the stream and using 16 both land on offered options.

End of Section 3 — Time, Work & Motion

You have worked drills 15–17 (3 of drills in this book). Any blank space below this line is deliberate — each section begins on a fresh page. Next: Section 4 — Data Interpretation & Equations, starting overleaf.

Data Interpretation & Equations — How to Read This Section

Tables, caselets, and paired quadratics — the marks are in reading only what the question asks for, then computing once.

Data Interpretation (Table). A small table and one question about it, and the question is always one of four: a column TOTAL, the DIFFERENCE between two named rows in one column, two columns COMBINED for a single row, or a column's AVERAGE. The table prints far more than any one question needs, so the skill is finding the named cells and ignoring the rest of the page.

WORKED EXAMPLE Branch / Men / Women — North 100 | 30, South 130 | 40, East 70 | 80. Total Men across all 3 branches? Only the Men column is in play: $100 + 130 + 70 = 300$. The Women column never enters it.

COMMON MISTAKE reading down the neighbouring column, or answering the question the table INVITES rather than the one printed. Underline the column name and the row names first — the arithmetic after that is a single addition.

Data Interpretation (Caselet). The same data as a table DI, hidden in a paragraph — and always in TWO steps. The paragraph splits a total (by a ratio or by a percentage), then asks for a share of ONE of the parts. Do the split on paper before you even look at what is being asked, and the question becomes a lookup.

WORKED EXAMPLE A store holds 48 items, split between imported and locally made in the ratio 5 : 7. Of the imported items, 25% are returned. How many imported items are returned? Step one, the split: 12 parts in all, so one part is 4 and imported = $5 \times 4 = 20$. Step two: 25% of 20 = 5. The 28 local items were never in the question.

COMMON MISTAKE taking that second percentage of the TOTAL instead of the part — 25% of 48 rather than of 20. The paragraph sets the trap by keeping both numbers in the same sentence.

Quadratic Equations. Two quadratics, one in x and one in y . Solve both, then compare EVERY root of one against every root of the other. The answer is a definite inequality only if the two sets do not overlap at all.

WORKED EXAMPLE $x^2 - 7x + 12 = 0$ and $y^2 - 9y + 20 = 0$. $x = 3$ or 4; $y = 4$ or 5. The ranges overlap — x can equal y at 4, and x can also be less. The relation cannot be determined.

COMMON MISTAKE comparing only the largest roots, or only the smallest. ' $x < y$ ' is true only when every single x is below every single y — one overlapping pair makes it indeterminate.

Section 4 Contents — Data Interpretation & Equations

Drills 18–20 of this section only. The full book index is at the front.

Drill 18 · Hard

Drill 20 · Medium

Drill 19 · Easy

Drill 18 **HARD** *Data Interpretation (Table)*[↑ Section index](#)**QUESTION****By how much does Correct differ between GA and Quant?****Given:**

1. One candidate's mock-test record, by section: (figures in questions)
2. Subject Attempted Correct
3. Quant 800 600
4. Reasoning 2200 600
5. English 600 400
6. GA 3200 1800

Options:

- A** 600
- B** 1198
- C** 2400
- D** 1200
- E** 1196

Target time: ~65s

⚠ **Common mistake:** "Differ" subtracts: 1800 and 600 within the Correct column. Their sum 2400 is on the option list, and so is the same difference read off the Attempted column (2400).

Drill 19 EASY *Data Interpretation (Caselet)*[↑ Section index](#)**QUESTION****How many express parcels are delivered next day?****Given:**

1. A depot dispatched 45 parcels, split between express and standard in the ratio 4 : 5.
2. Of the express parcels, 25% are delivered next day.

Options:

- A** 5
- B** 20
- C** 2
- D** 4
- E** 15

Target time: ~40s

⚠ **Common mistake:** The 25% applies to the express group (20), not to all 45 parcels — applying it to the total is the single commonest caselet slip, and it is on the option list.

Drill 20 MEDIUM *Quadratic Equations*[↑ Section index](#)Solve both equations and state the relation between x and y .**Given:**

1. I. $x^2 - 22x + 120 = 0$
2. II. $y^2 - 9y + 20 = 0$

Options:

- A** $x > y$
- B** $x < y$
- C** $x \geq y$
- D** $x \leq y$
- E** $x = y$ or no relation can be established

Target time: ~40s

⚠ **Common mistake:** comparing only the largest roots, or only the smallest. ' $x < y$ ' is true only when every single x is below every single y — one overlapping pair makes it indeterminate.

End of Section 4 — Data Interpretation & Equations

You have worked drills 18–20 (3 of drills in this book). Any blank space below this line is deliberate — each section begins on a fresh page. Next: Section 5 — Counting & Measurement, starting overleaf.

Counting & Measurement — How to Read This Section

Arrangements, chances, and shapes — the topics where the setup is the whole battle and the arithmetic afterwards is trivial.

Permutation & Combination. Ask one question first: does ORDER matter? A committee, a handshake or a selection is a set, so it is a combination; a ranking, a seating or a code is an arrangement, so it is a permutation. Then split any 'at least' condition into cases and add them.

WORKED EXAMPLE From 5 men and 4 women, choose a committee of 3 with at least 2 women. Split by cases:
exactly 2 women = $C(4,2) \times C(5,1) = 6 \times 5 = 30$; exactly 3 women = $C(4,3) = 4$. $30 + 4 = 34$.

COMMON MISTAKE using a permutation where order does not matter — a committee of A, B, C is the same committee however you list it, so it is counted once, not six times.

Probability. Favourable outcomes over total outcomes, with both counted the same way. Most drills are a single draw — one ball from a bag, one bulb from a box, one card from a pack of 52, or the sum on two dice out of 36 — so the work is counting the favourable cases exactly. Only when two are drawn WITHOUT replacement does the second draw see a changed bag.

WORKED EXAMPLE Two fair dice are rolled together. P(the sum is exactly 5)? Total outcomes = $6 \times 6 = 36$. A sum of 5 happens as 1+4, 2+3, 3+2, 4+1 — four ordered pairs. $4 \div 36 = 1/9$.

COMMON MISTAKE counting 1+4 and 4+1 as one outcome. The 36 in the denominator treats the two dice as distinguishable, so the numerator has to as well — otherwise the two halves of the fraction are counting different experiments.

Mensuration. Raw formulas on the standard shapes, and only a handful of them: area and perimeter of a rectangle and a square, the volume of a cuboidal tank, and the area and circumference of a circle. Circle questions hand you $\pi = 22/7$, which is itself a hint — the radius will be a multiple of 7, so the 7 always cancels.

WORKED EXAMPLE A circular park has a radius of 21 m. (Take $\pi = 22/7$.) What is its circumference? $C = 2\pi r = 2 \times 22/7 \times 21$, and 7 goes into 21 three times — so this is $2 \times 22 \times 3 = 132$ m.

COMMON MISTAKE answering the OTHER formula for the same shape — the area when it asked for the perimeter, the circumference when it asked for the area, the surface area when it asked for the volume. That answer is always sitting in the options. Read the last four words of the question before you touch a number.

Section 5 Contents — Counting & Measurement

Drills 21–23 of this section only. The full book index is at the front.

Drill 21 · Hard

Drill 23 · Medium

Drill 22 · Easy

Drill 21 **HARD** Mensuration[↑ Section index](#)**QUESTION****What is its circumference?****Given:****A circular park has a radius of 84 m. (Take $\pi = 22/7$.)****Options:**

- A** 522
- B** 264
- C** 22176
- D** 528
- E** 168

Target time: ~60s

⚠ **Common mistake:** Circumference is $2\pi r$, linear in 84; squaring the radius gives the AREA 22176 m², which is on the option list. The "2" is easy to drop — half of 528 is 264, also offered.

Drill 22 **EASY** Permutation & Combination[↑ Section index](#)**QUESTION****In how many different ways can they be arranged?****Given:****6 students are to stand in a single row for a photograph.****Options:**

- A** 36
- B** 30
- C** 720
- D** 120
- E** 12

Target time: ~45s

⚠ **Common mistake:** using a permutation where order does not matter — a committee of A, B, C is the same committee however you list it, so it is counted once, not six times.

Drill 23**MEDIUM**

Probability

[↑ Section index](#)**QUESTION****What is the probability that it is NOT defective?****Given:**

1. A box contains 5 defective and 4 good bulbs.
2. One bulb is picked at random.

Options:

- A** $2/5$
- B** $4/5$
- C** $1/9$
- D** $5/9$
- E** $4/9$

Target time: ~55s

⚠ **Common mistake:** counting 1+4 and 4+1 as one outcome. The 36 in the denominator treats the two dice as distinguishable, so the numerator has to as well — otherwise the two halves of the fraction are counting different experiments.

End of Section 5 — Counting & Measurement

You have worked drills 21–23 (3 of drills in this book). Any blank space below this line is deliberate — each section begins on a fresh page. Next: the Solutions — the answer and the full worked method for every puzzle, in the same numbering.

Solutions

Worked solutions for every drill — the answer plus the engine's own step-by-step working. Attempt the drill against its target time first; then read the working even if you were right, because the key's method is usually the shorter one.

Solution 1 EASY Speed Arithmetic

Q. Solve in your head — beat the clock.

Answer: A) 72

Why:

1. Compute directly: $8 \times 9 = 72$.

Solution 2 MEDIUM Simplification (BODMAS)

Q. Simplify — order of operations wins.

Answer: B) 82

Why:

1. $\times$ and $\div$ first: $4 \times 4 = 16$; $45 \div 9 = 5$.
2. Then left to right: $71 + 16 - 5 = 82$.

Solution 3 HARD Squares, Cubes & Roots

Q. Powers and roots — instant recall.

Answer: C) 7

Why:

1. $7^3 = 343$, so $\sqrt[3]{343} = 7$. The last digit of a cube pins the root's last digit uniquely.

Solution 4 EASY Vedic Squaring

Q. Squaring a number ending in 5 — beat the clock.

Answer: D) 3025

Why:

1. Ends in 5 — Ekadhikena: take 5, multiply by one more: $5 \times 6 = 30$.
2. Append 25: $30 \mid 25 \rightarrow 3025$.
3. Answer: 3025.

Solution 5 MEDIUM Crosswise Multiplication

Q. Crosswise multiplication — beat the clock.

Answer: C) 5292

Why:

1. Units: $3 \times 4 = 12$.
2. Cross: $6 \times 4 + 3 \times 8 = 48$; tens: $6 \times 8 = 48$ — stack with carries.
3. $4800 + 480 + 12 = 5292$.
4. Answer: 5292.

Solution 6 **HARD** *Nikhilam*

Q. Nikhilam multiplication — beat the clock.

Answer: C) 11872

Why:

1. Choose the base (nearest power of 10): 100.
2. Distance from base: 106 is 6 above 100, 112 is 12 above 100.
3. Cross-add: $106 + 12 = 118$ (same as $112 + 6$).
4. Multiply the distances: $6 \times 12 = 72$.
5. Base 100 has 2 zeros, so the right-hand part must occupy exactly 2 digits.
6. 72 fits, padded to 2 digits $\rightarrow 72$.
7. Left part: 118. Right part: 72.
8. Write them side by side: $118 | 72 = 11872$.

Solution 7 **EASY** *Complementary Products*

Q. Units sum to 10, same lead — beat the clock.

Answer: E) 7216

Why:

1. Units $8 + 2 = 10$ with the same lead 8 — Antyayordashake applies.
2. Left: $8 \times 9 = 72$. Right: $8 \times 2 = 16$ (always two digits).
3. $72 | 16 \rightarrow 7216$.
4. Answer: 7216.

Solution 8 **MEDIUM** *Difference of Squares*

Q. Multiply via the midpoint — beat the clock.

Answer: E) 836

Why:

1. 22 and 38 straddle 30 by 8 — Sankalana: $a \times b = m^2 - d^2$.
2. $30^2 - 8^2 = 900 - 64 = 836$.
3. Answer: 836.

Solution 9 **HARD** *Paravartya Division*

Q. Dividing by 11 — beat the clock.

Answer: B) 9005

Why:

1. $99055 \div 11$ — Paravartya: $11 = 10 + 1$, so the flag digit is 1; negate it to -1 .
2. Bring down the first digit as-is: 9 (first quotient digit).
3. $9 + (-1 \times 9) = 0$ (next quotient digit).
4. $0 + (-1 \times 0) = 0$ (next quotient digit).
5. $5 + (-1 \times 0) = 5$ (next quotient digit).
6. Last digit gives the remainder the same way: $5 + (-1 \times 5) = 0$.
7. Quotient 9005, remainder 0 — check: $9005 \times 11 + 0 = 99055$.
8. Answer: 9005.

Solution 10 **EASY** *Anurupyena Multiplication*

Q. Multiplying near a working base — beat the clock.

Answer: E) 2208

Why:

1. Deviations from 50: $48 \rightarrow -2$, $46 \rightarrow -4$.
2. Cross-add for the raw left part: $48 - 4 = 44$.
3. Rescale for the working base: 50 is $100 \div 2$, so divide by 2: $44 \div 2 = 22$.
4. Right part — multiply the deviations: $(-2) \times (-4) = 8$, padded to two digits: 08.
5. Join them: $22 \mid 08 \rightarrow 2208$.
6. Answer: 2208.

Solution 11 **MEDIUM** *Percentages*

Q. Percentage drill — no paper.

Answer: E) 75

Why:

1. The change is $44 - 11 = 33$.
2. Percentage change is always measured on the ORIGINAL value: $33 / 44 \times 100 = 75\%$.

Solution 12 **HARD** *Profit & Loss*

Q. What was the cost price?

Answer: E) 8000

Why:

1. A profit of 5% means $SP = 105\%$ of CP.

2. So CP = ₹8400 × 100 ÷ 105 = ₹8000 — the percentage is always on COST price, never on the selling price.

3. Answer: 8000.

Solution 13 **EASY** Simple & Compound Interest

Q. What is the simple interest earned?

Answer: A) 4000

Why:

1. Simple interest = $P \times R \times T \div 100 = ₹10000 \times 20 \times 2 \div 100 = ₹4000$.
2. Each year earns the same ₹2000 under simple interest.
3. Answer: 4000.

Solution 14 **MEDIUM** Averages

Q. Averages — total and mean, no paper.

Answer: A) 39

Why:

1. All 6 together: $6 \times 50 = 300$.
2. Subtract the 5 you were given: $300 - 261 = 39$. The remaining number is 39.

Solution 15 **HARD** Time & Work

Q. Working together, in how many days will they finish it?

Answer: B) 4

Why:

1. Take the total work as 12 units (the LCM of 6 and 12).
2. Anil does 2 units/day, Bhavna does 1 units/day — add the RATES, never the days.
3. Together they do 3 units/day, so the job takes $12 \div 3 = 4$ days.
4. Answer: 4 days.

Solution 16 **EASY** Time, Speed & Distance

Q. How far does it travel?

Answer: A) 90

Why:

1. Distance = speed × time = $30 \times 3 = 90$ km.
2. Answer: 90 km.

Solution 17 MEDIUM Boats & Streams

Q. How far does it go?

Answer: A) 55

Why:

1. Upstream speed = $16 - 5 = 11$ km/h.
2. Distance = $11 \times 5 = 55$ km — the stream must be applied before multiplying.
3. Answer: 55 km.

Solution 18 HARD Data Interpretation (Table)

Q. By how much does Correct differ between GA and Quant?

Answer: D) 1200

Why:

1. GA Correct = 1800, Quant Correct = 600.
2. Difference = 1200 — subtract within one column.
3. Answer: 1200.

Solution 19 EASY Data Interpretation (Caselet)

Q. How many express parcels are delivered next day?

Answer: A) 5

Why:

1. The ratio 4 : 5 has 9 parts, so one part = $45 \div 9 = 5$.
2. Express parcels = $4 \times 5 = 20$.
3. 25% of 20 = 5 — take the percentage on the PART, never the total.
4. Answer: 5.

Solution 20 MEDIUM Quadratic Equations

Q. Solve both equations and state the relation between x and y.

Answer: A) $x > y$

Why:

1. Factorise I: $x^2 - 22x + 120 = 0 \rightarrow (x - 10)(x - 12) = 0$, so $x = 10$ and 12.
2. Factorise II: $y^2 - 9y + 20 = 0 \rightarrow (y - 4)(y - 5) = 0$, so $y = 4$ and 5.
3. Compare every x against every y — one pair is never enough.
4. The smallest x (10) still beats the largest y (5), so $x > y$.

Solution 21 **HARD** *Mensuration*

Q. What is its circumference?

Answer: D) 528

Why:

1. Circumference = $2\pi r = 2 \times (22/7) \times 84 = 528$ m — circumference uses r , area uses r^2 .
2. Answer: 528 m.

Solution 22 **EASY** *Permutation & Combination*

Q. In how many different ways can they be arranged?

Answer: C) 720

Why:

1. Each position is filled in turn: 6 choices, then 5, and so on down to 1.
2. Arrangements = $6! = 6 \times 5 \times 4 \times 3 \times 2 \times 1 = 720$.
3. Answer: 720.

Solution 23 **MEDIUM** *Probability*

Q. What is the probability that it is NOT defective?

Answer: E) $4/9$

Why:

1. $P(\text{not defective}) = 1 - P(\text{defective}) = 1 - 5/9$.
2. That is the 4 good bulbs out of 9: $4/9$.
3. Answer: $4/9$.

YOU'VE FINISHED THE BOOK

Keep the clock running

You've drilled every type and read the working — now keep the speed. ComputePrep serves a fresh timed set **every single day**, free, at computeprep.com: the same target times, a streak, and unlimited practice on whichever type you keep missing on time.

Start Today's Drills →

Quant is half the paper. Reasoning is the other half.

An IBPS / SBI prelims paper is 100 questions — 35 Quantitative Aptitude, 35 Reasoning Ability, 30 English — with a fixed clock on each section; SSC CGL Tier-I gives Quant and Reasoning 25 questions apiece. Speed on the calculation side is what this book trains. On the reasoning side the biggest single block is usually the **puzzle / seating-arrangement set** — one arrangement, five questions, and the same skill: fewer steps under time. **Deduce**, our sister app, is that block as a daily game — one server-checked logic puzzle a day, three attempts, a streak and a leaderboard. Same engine discipline as this book: every puzzle is generated and verified to have exactly one answer before it is served.

⚡ Today's drills — free, no signup

computeprep.com — a fresh timed set daily across all 23 types, plus unlimited practice on your weakest one.

▶ Play Deduce — today's reasoning puzzle, free

deducepuzzles.com — seating, floor and scheduling puzzles tuned to your exam (Banking, SSC, Railway, UPSC CSAT, State PSC, GRE/GMAT), one a day, no signup to play.

★ Pro — one plan, both apps

₹199/month or ₹1,499/year, 7 days free. Unlimited drills + the Ganak coach on ComputePrep; past-puzzle archive, unlimited practice + the Saarathi coach on Deduce. computeprep.com/pro

■ Deduce eBooks — reasoning, same format as this book

100 Reasoning Puzzles per exam (Banking · SSC · Railway · UPSC CSAT · State PSC · GRE/GMAT), ₹299 each, or 150 Reasoning Puzzles for Bank Exams from ₹99; free 20-puzzle sample. deducepuzzles.com/ebook

◆ The full Speed Book · ₹199 once

150 drills instead of 23, every type at three difficulties, full worked key. computeprep.com/speed-book

☰ Free Vedic Speed Kit (PDF)

The shortcuts on one printable sheet — squaring, base multiplication, $\times 11$. computeprep.com/speed-kit

FOLLOW ALONG

t.me/computeprep
youtube.com/@computeprep
instagram.com/computeprep
t.me/deducepuzzles
youtube.com/@deducepuzzles
instagram.com/deducepuzzles

A product of **Launchship IT & Media Services Pvt. Ltd.** · computeprep.com · deducepuzzles.com · Feedback or an error to report: hello@computeprep.com

Free Sample · © 2026 Launchship IT & Media Services Pvt. Ltd. All rights reserved. This free sample may be shared and forwarded freely. Every drill is generated by the ComputePrep engine, and its answer and worked solution are computed by that same engine. ComputePrep and Deduce are independent practice tools, not affiliated with any exam body; exam patterns quoted are as published at the time of writing.

FROM THE SAME STUDIO · LAUNCHSHIP

Now train the other half of the paper: reasoning

Deduce — The reasoning half of the paper — one server-checked logic puzzle a day.

A coach on each side — one Pro plan unlocks both



COMPUTEPREP · QUANT COACH · HERE

Ganak

After a drill he explains the fastest route — the Vedic shortcut, and where your method lost the seconds. Everyone meets him on their first drill. **With Pro: on every drill verdict, 120 turns a day.**



DEDUCE · REASONING COACH

Saarathi

Ask “why is that the answer?” — he walks the deduction step by step, tracks the puzzle types that cost you marks and tells you what to drill next. **With Pro: 120 coach turns a day, on any puzzle.**

The Deduce app — free, every day

- One fresh puzzle every day (IST) — seating, floor, box, scheduling, ranking, inequality, syllogism, blood relations, direction, coding and more
- Every puzzle is engine-generated and proven to have exactly ONE solution before you see it — never hand-written, never ambiguous
- 3 attempts, server-timed; a streak, a daily leaderboard and a shareable result card
- Pick your exam — Banking, SSC, Railway, UPSC CSAT, State PSC or GRE/GMAT — and the daily tunes its topic mix to that paper
- Private groups: a leaderboard with your own study circle
- The morning puzzle on Telegram, free — t.me/deducepuzzles

Deduce with Pro

- Past Puzzles — replay the whole archive
- Unlimited adaptive practice on the types you keep missing
- Saarathi, the reasoning coach — 120 turns a day, on any puzzle
- 3 board hints per daily (instead of 1) when you're stuck

★ Pro — one plan, both apps

₹199 / month or ₹1,499 / year · 7 days free

One subscription unlocks Pro on Deduce AND ComputePrep — reasoning and calculation, the two halves of the paper, for the price of one.

The Deduce eBooks — what to expect

■ 100 Reasoning Puzzles — one edition per exam · ₹299 each

Banking (IBPS/SBI) · SSC · Railway · UPSC CSAT · State PSC · GRE/GMAT. Each is 100 solver-validated puzzles weighted to that exam's own topic mix, Easy → Hard, sectioned by type, with a type-by-type “how to read this arrangement” primer and a clue-by-clue worked solution for every puzzle.

■ 150 Reasoning Puzzles for Bank Exams · from ₹99

The complete banking-reasoning syllabus in one book — every puzzle type, three difficulty levels, full worked solutions. All seven books together: ₹1,499.

📖 Free download: 20 reasoning puzzles — free PDF

A 20-puzzle sample from the Deduce books — plus a separate free worked-solutions PDF and a fresh puzzle PDF pack on the free-puzzles page. Forward it to anyone prepping with you.

deducepuzzles.com/ebook/deduce-20-puzzles-sample.pdf

Or start on the app, free: **deducepuzzles.com** · More free PDFs: **deducepuzzles.com/free-puzzles**



DEDUCE

t.me/deducepuzzles

youtube.com/@deducepuzzles

instagram.com/deducepuzzles

COMPUTEPREP

t.me/computeprep

youtube.com/@computeprep

instagram.com/computeprep

Deduce and ComputePrep are products of Launchship IT & Media Services Pvt. Ltd. — **deducepuzzles.com** · **computeprep.com**. Prices in INR as published at the time of writing. Independent practice tools, not affiliated with IBPS, SBI, SSC, RRB, UPSC, GRE/GMAT or any state PSC.